

HEDIS Provider Guide: Chlamydia Screening in Women (CHL)

Measure Description

Women 16 to 24 years of age who were identified as sexually active and who had at least one chlamydia test during the measurement year.

Members identified as sexually active:

- Sexually Transmitted Infection (STI) screening
- Contraceptive prescribed
- Pregnancy test completed

Exclusions: Members in hospice.

Using Correct Billing Codes

Description	Codes
Chlamydia Screening	<p>CPT: 87110, 87270, 87320, 87490-87492, 87810, 0353U</p> <p>LOINC: 14463-4, 14464-2, 14465-9, 14467-5, 14474-1, 14513-6, 16600-9, 21190-4, 21191-2, 21613-5, 23838-6, 31775-0, 34710-4, 42931-6, 43304-5, 43404-3, 44806-8, 44807-6, 45068-4, 45069-2, 45072-6, 45073-4, 45075-9, 45084-1, 45089-0, 45090-8, 45091-6, 45093-2, 45095-7, 4993-2, 50387-0, 53925-4, 53926-2, 57287-5, 6353-7, 6356-0, 6357-8, 80360-1, 80361-9, 80362-7, 80363-5, 80364-3, 80365-0, 80367-6, 82306-2, 87949-4, 87950-2, 88221-7, 89648-0, 91860-7, 91873-0</p>

How to Improve HEDIS® Scores

- Perform chlamydia screening every year on every 16 to 24-year-old female identified as sexually active (use any visit opportunity).
- Add chlamydia screening as a standard lab for women 16 to 24 years old. Use adolescent well-care exams and well woman exams for this purpose.
- Remember that chlamydia screening can be performed through a urine test. Offer this option for your patients.
- Request a focused list of members who need chlamydia screening from your Blue Shield Promise Quality Program Manager.

HEDIS® is a registered trademark for the National Committee of Quality Assurance (NCQA)